FTUV-15-09-25
IFIC-15-50

On the tensor-to-scalar ratio in large single-field inflation models

Gabriela Barenboim* and Wan-Il Park†
Departament de Física Teòrica and IFIC, Universitat de València-CSIC, E-46100, Burjassot, Spain
(Dated: September 25, 2015)

We show that generically the tensor-to-scalar ratio in large single-field inflation scenarios is bounded to be larger than $\mathcal{O}(10^{-3})$ for the spectral index in the range favored by observations.

arXiv:1509.07132v1 [astro-ph.CO] 23 Sep 2015

INTRODUCTION

In scenarios of inflation, a necessary step to explain various features of the present universe, the number of e-foldings (N_e) is conventionally taken to be around 50-60 depending on the cosmological history after inflation. The lower bound of N_e can be pushed down as long as the reheating temperature after inflation is high enough to provide the relevant physics afterwards (e.g., baryogenesis, production of dark matter, a successful Big-Bang nucleosynthesis). However, the upper bound is saturated at about 60, since the upper bound of the tensor-to-scalar ratio ($r_T < 0.1$ [1]) coming from the non-observations of the tensor mode perturbations, puts an upper bound on the energy scale of inflation *i.e.,* $V_*^{1/4} \lesssim 2 \times 10^{16}\,\mathrm{GeV}$.

In conventional Einstein gravity, the tensor-to-scalar ratio is simply proportional to the slow-roll parameter associated with the slop of the inflaton potential. For small field inflation in which the excursion scale of the inflaton can be lowered down far below Planck scale, r_T can be extermely small. However, in large-field inflation scenarios, this can not be the case since the e-foldings are likely to turn out too large unless the potential has an unphysical sudden change. Hence, r_T is likely to be lower-bounded in such scenarios.

In this work, we show that the e-foldings required to match observations set a lower bound of r_T at around $\mathcal{O}(10^{-3})$ in most of realistic models of large single-field inflation. Our result is consistent with a recent work, Ref. [2], although the approach is different.

INFLATION AND THE NUMBER OF e-FOLDINGS FOR A GIVEN SCALE

In slow-roll inflation scenario with a potential V of a single scalar field (playing the role of inflaton), the slow-roll parameters are defined as

$$\epsilon \equiv \frac{1}{2}\left|\frac{M_{\rm P}V'}{V}\right|^2, \quad \eta \equiv \frac{M_{\rm P}^2 V''}{V}, \quad \xi^2 \equiv \frac{M_{\rm P}^4 V'V''}{V^2} \tag{1}$$

where $M_{\rm P} = 2.4 \times 10^{18}\,\mathrm{GeV}$ is the reduced Planck mass, and '$\prime$' represents derivative with respect to the inflaton field. The power spectrum of scalar and tensor mode perturbations are given by

$$P_{\mathcal{R}} = \left(\frac{H}{2\pi}\right)^2 \frac{1}{2\epsilon M_{\rm P}^2}, \quad P_T = \left(\frac{H}{2\pi}\right)^2 \frac{8}{M_{\rm P}^2} \tag{2}$$

respectively, giving a tensor-to-scalar ratio

$$r_T \equiv \frac{P_T}{P_{\mathcal{R}}} = 16\epsilon \tag{3}$$

The spectral indices of these modes are

$$n_s = 1 - 6\epsilon + 2\eta, \quad n_T = -2\epsilon \tag{4}$$

Note that $n_T = -r_T/8$, a consistency relation of inflation in Einstein gravity. For inflation to end smoothly, ϵ should depend on the scale; although η can be constant. The spectral running, $\alpha \equiv dn_s/d\ln k$, is given by

$$\alpha = 16\epsilon\eta - 24\epsilon^2 - 2\xi^2 \tag{5}$$

The number of e-foldings generated until the end of inflation, since a given cosmological scale (denoted with a subscript '$_*$') exits the horizon during inflation, is given by

$$N_{e,*}^{\rm th} \equiv \int_*^e H dt \simeq -\frac{1}{M_{\rm P}^2}\int_*^e d\phi \frac{V}{V'} \tag{6}$$

with ϕ being the inflaton field. For a given model of inflation (i.e., a specific form of potential V), the observed n_s fixes ϕ_* and hence $N_{e,*}^{\rm th}$ is also fixed. Meanwhile, in order for inflation to solve the smoothness problem of Big-Bang cosmology, any two points within our Hubble patch had to be in casual contact at some stage before the end of inflation. That requires a minimum amount of e-foldings, that we will denote $N_{e,*}^{\rm obs}$. In order for an inflation scenario to work, $N_{e,*}^{\rm th} = N_{e,*}^{\rm obs}$ is required, and this constrains the possible models of inflation. If there were no more entropy production after the reheating following inflation, the e-foldings associated with the Planck pivot scale ($k_* = 0.05\mathrm{Mpc}^{-1}$) are given by [3]

$$\begin{aligned} N_{e,*}^{\rm obs} &= \frac{1}{3}\ln\left[\frac{2\sqrt{6}\pi^3}{3}\frac{s_0}{k_*^3}\left(\frac{V_*}{V_e}\frac{V_e^{\frac{1}{4}}}{M_{\rm P}}\right)^3\left(\frac{T_{\rm d}}{V_e^{\frac{1}{4}}}\right)\right] \\ &\simeq 57.87 - \frac{1}{3}\ln\left[\left(\frac{V_e}{V_*}\right)^3\left(\frac{10^{16}\,\mathrm{GeV}}{V_e^{\frac{1}{4}}}\right)^3\frac{V_e^{\frac{1}{4}}}{T_{\rm d}}\right] \end{aligned} \tag{7}$$

where s_0 is the present entropy density

$$s_0 = \frac{2\pi^2}{45} g_{*s0} T_0^3 \tag{8}$$

with $g_{*s0} = 3.9091$ being the number of relativistic degrees of freedom and $T_0 = 2.726\text{K}$ the CMB temperature at the present universe. The subscript '$_e$' in the right-hand side denotes the value at the end of inflation, and T_d is the decay temperature of the inflaton. The current upper-bound on the tensor-to-scalar ratio $r_T < 0.1$ [1] implies

$$V_*^{1/4} \lesssim 2 \times 10^{16}\,\text{GeV} \tag{9}$$

and the reheating temperature is bounded as

$$1\,\text{MeV} \lesssim T_\text{d} \lesssim V_*^{1/4} \tag{10}$$

Then, setting $V_* = V_e$, one finds

$$N_{e,*}^\text{obs} \lesssim 58.56 \tag{11}$$

The lower bound of $N_{e,*}^\text{obs}$ depends on the post-inflation cosmology as well as V_* and T_d. In particular, there can be an $\mathcal{O}(10)$ additional contribution to the e-foldings of inflation from, for example, thermal inflation [4, 5]. In this case, the required e-foldings from primordial inflation are reduced by that amount. In principle, there can be multiple stages of thermal inflation, providing a few tens of e-foldings. However, thermal inflation with more than two stages is non-trivial to realize (or not so realistic), and too much extra e-foldings may cause some inconsistency with observations of small scale structure (see for example [6]). In realistic models, we expect extra e-foldings of around 20 or less [7–11]. Hence, in this letter we take $\Delta N_e = 15$ as the plausible maximal extra e-foldings with $T_\text{d} = V_*^{1/4}$ for simplicity in the case where thermal inflation is considered.

The number of e-foldings in Eq. (6) can be separated into contributions from slow-roll and non-slow-roll (or fast-roll) regions, denoted respectively as $\Delta N_{e,\text{sr}}$ and $\Delta N_{e,\text{fr}}$, i.e.,

$$N_{e,*}^\text{th} = \Delta N_{e,\text{sr}} + \Delta N_{e,\text{fr}} \tag{12}$$

where

$$N_{e,\text{sr}} = \int_*^\times \frac{d\phi/M_\text{P}}{\sqrt{2\epsilon}}, \quad N_{e,\text{fr}} = \int_\times^e \frac{d\phi/M_\text{P}}{\sqrt{2\epsilon}} \tag{13}$$

The value of $\phi_\times$ for which the slow-roll approximation breaks down depends on the potential and is rather ambiguous. However, we can take $\epsilon_\times \equiv \epsilon(\phi_\times) \sim \mathcal{O}(0.1)$, and in this case

$$N_{e,\text{fr}} < \frac{\Delta\phi/M_\text{P}}{\epsilon_\times} \simeq 2.24 \left(\frac{0.1}{\epsilon_\times}\right)^{1/2} \frac{\Delta\phi}{M_\text{P}} \tag{14}$$

where $\Delta\phi \equiv |\phi_e - \phi_\times|$ with ϕ_e being the value of the inflaton at the end of inflation. In large-field models, generically, $\Delta\phi \gtrsim M_\text{P} \sim 1$, being the precise value model dependent. Hence, as a simple conservative constraint, we require

$$\Delta N_{e,\text{sr}} < N_{e,*}^\text{th} - \Delta N_e \tag{15}$$

with $\Delta N_e = 0, 15$ depending on the existence of thermal inflation.

CLASSES OF POTENTIALS

There are numerous different shapes of potentials for slow-roll inflation. However, in the region of field values where the slow-roll approximation is still valid, inflaton potentials generally fall into one of the following forms,

- Chaotic monomial: $V_\text{ch} = V_0 x^p$
- Inverse-Hilltop: $V_\text{iht} = V_0\,(1 - 1/x^p + \ldots)$
- Starobinsky-like: $V_\text{st} = V_0\,(1 - e^{-x} + \ldots)$
- Hilltop: $V_\text{ht} = V_0\,(1 - x^p + \ldots)$

where $x = \phi/\mu$ with μ being a scale characterizing the end of inflation, and p is assumed to be positive definite. Note that, even if the potentials listed above are approximated froms, it is clear that they all are significantly changed as $x \to 1$. Hence, unless the potentials have non-trivial complications around the end of inflation, we expect that inflation ends at $x \sim 1$. In the following subsections, we show the approximate forms of slow-roll parameters and $N_{e,*}^\text{th}$ or $N_{e,\text{sr}}$ (depending on its relevance) which are expected to be valid as long as ϵ and η are much smaller than unity in the region where $V \approx V_0$. These approximate expressions are useful to get an idea of the parametric dependences of the relevant quantities. The discrepancies one would get when considereing a complete potential are minor and do not change our argument.

Chaotic monomial

In this case, the slow-roll parameters take the (exact) form

$$\epsilon = \frac{p^2}{2x^2}\left(\frac{M_\text{P}}{\mu}\right)^2, \quad \eta = \frac{p(p-1)}{x^2}\left(\frac{M_\text{P}}{\mu}\right)^2 \tag{16}$$

giving $\eta = 2(p-1)\epsilon/p$, and from Eq. (4)

$$\epsilon_* = \left(\frac{p}{p+2}\right)\frac{1-n_s}{2} \tag{17}$$

which does not depend on μ. The e-foldings are

$$N_{e,*}^{\rm th} = \frac{p}{4}\left(\frac{1}{\epsilon_*} - 1\right) \simeq \frac{p+2}{2(1-n_s)} \tag{18}$$

It is clear that, as p becomes larger than 1, η becomes larger than zero, and similarly ϵ. Also, only $p \lesssim 4$ is allowed, othewise too much e-foldings are expected. As can be seen from Eq. (17), ϵ can be lowered down by taking a small p. However, in large field scenarios in which $x_* > 1$ with $\mu \geq M_{\rm P}$, we are constrained to have $p \gtrsim \mathcal{O}(10^{-2})$. Hence, r_T is lower-bounded at $\mathcal{O}(10^{-3})$.

Inverse-Hilltop potential

For $V \approx V_0$, the slow-roll parameters can be approximated as

$$\epsilon \approx \frac{p^2}{2x^{2(p+1)}}\left(\frac{M_{\rm P}}{\mu}\right)^2, \quad \eta \approx -\frac{p(p+1)}{x^{p+2}}\left(\frac{M_{\rm P}}{\mu}\right)^2 \tag{19}$$

From Eq. (19), we find

$$\eta = -\frac{2(p+1)}{p}\left[\frac{2}{p^2(M_{\rm P}/\mu)^2}\right]^{q-1}\epsilon^q \tag{20}$$

where

$$\frac{1}{2} < q \equiv \frac{p+2}{2(p+1)} < 1 \tag{21}$$

Using Eq. (4) with Eq. (20), one can find ϵ_*, at least numerically, as a function of p and n_s for a given μ. Note that $\eta/\epsilon = -2x^p(p+1)/p$ in this potential, which means that for a given η, as p goes away from a value around one (or μ is lowered down), ϵ decreases. The e-foldings for slow-roll regime are

$$N_{e,\rm sr} \approx \frac{x_*^{p+2} - x_\times^{p+2}}{p(p+2)}\left(\frac{\mu}{M_{\rm P}}\right)^2 \approx \left(\frac{1}{\eta_\times} - \frac{1}{\eta_*}\right)\left(\frac{p+1}{p+2}\right) \tag{22}$$

where $\eta_\times$ is found from Eq. (20) with $\epsilon_\times = 0.1$.

We can explore the limiting cases of p leading to small ϵ_*. When $p \to 0$, one finds

$$\epsilon_* \simeq \frac{p}{4}(1-n_s), \; N_{e,\rm sr} \simeq \frac{1}{1-n_s} \tag{23}$$

Hence, taking a small p, one can lower down ϵ_*, but $N_{e,\rm sr} \gtrsim 50$ requires $n_s \gtrsim 0.98$ which is out of the 2-σ region. On the other hand, if $p \to \infty$,

$$\epsilon_*^{1/2} \simeq -\frac{\gamma}{6}\left(1 - \sqrt{1 + \frac{6(1-n_s)}{\gamma^2}}\right), \; N_{e,\rm sr} \simeq \frac{2}{1-n_s} \tag{24}$$

with $\gamma \equiv \sqrt{2}p(M_{\rm P}/\mu)$. It may be natural and tempting to set μ to be the Planck scale. However, in principle μ can be much larger than $M_{\rm P}$ although it may need some fine-tuning or non-trivial (or unnatural) realizations.

Starobinsky-like

In this case, one finds

$$\epsilon \approx \frac{e^{-2x}}{2}\left(\frac{M_{\rm P}}{\mu}\right)^2, \quad \eta \approx -e^{-x}\left(\frac{M_{\rm P}}{\mu}\right)^2 \tag{25}$$

Note that, as μ is increased, ϵ_* matching observations increases as well. So, we take $\mu = M_{\rm P}$ to see the smallest allowed ϵ_*. From Eq. (25), one finds

$$\eta = -\sqrt{2}\epsilon^{1/2} \tag{26}$$

leading to

$$\epsilon_*^{1/2} = -\frac{\sqrt{2}}{6}\left(1 - \sqrt{1+3(1-n_s)}\right) \tag{27}$$

which is the same as $\epsilon_*^{1/2}$ in Eq. (24) with $\mu/M_{\rm P} = p$. The the number of e-foldings for slow-roll regime are

$$N_{e,\rm sr} \approx \frac{1}{\eta_\times} - \frac{1}{\eta_*} \tag{28}$$

where $\eta_\times$ is obtained from Eq. (26) with $\epsilon_\times = 0.1$.

Hilltop potential

In this case, for $0 < p < 1$ inflaton should be located at a particular region in $0 < x < 1$ as the initial condition, and this is non-trivial to realize. Also, as p increases, the ϵ_* needed to match observation becomes smaller. So, we consider only $p \geq 2$ in order to avoid irrelevant complications. Slow-roll parameters are given as

$$\epsilon \approx \frac{p^2 x^{2(p-1)}}{2}\left(\frac{M_{\rm P}}{\mu}\right)^2, \quad \eta \approx -p(p-1)x^{p-2}\left(\frac{M_{\rm P}}{\mu}\right)^2 \tag{29}$$

Note that for $p = 2$, η is a constant depending on μ exclusively, and

$$\frac{\mu}{M_{\rm P}} = \frac{2}{\sqrt{1-n_s-6\epsilon_*}} \geq \frac{2}{\sqrt{1-n_s}} \sim \mathcal{O}(10) \tag{30}$$

For $p \neq 1, 2$, η can be found in terms of ϵ from Eqs. (20) and (21) with $p \to -p$. The number of e-foldings is

$$N_{e,\rm sr} \approx \begin{cases} \frac{1}{2\eta}\ln\left(\frac{\epsilon_\times}{\epsilon_*}\right) & : \; p = 2 \\ \left(\frac{1}{\eta_\times} - \frac{1}{\eta_*}\right)\left(\frac{p-1}{p-2}\right) & : \; p \neq 1, 2 \end{cases} \tag{31}$$

Similarly to the case of $V_{\rm iht}$, as μ decreases, ϵ_* decreases, and we consider $\mu/M_{\rm P} \geq 1$ here too. Note that, if $\mu/M_{\rm P} = p$, as $p \to \infty$, ϵ_* collapses to the one in Eq. (27).



FIG. 1: The lower bound of r_T (gray line(s)) as a function of n_s. Green regions are 1, 2, 3-σ bands of n_s from Planck observations (Planck TT+lowP) [12]. Red lines are obtained by imposing the constraint on the e-foldings, Eq. (15). Red dashed lines are for inverse-Hilltop potential with $p = 0.1, 4$ from right to left. Red dotdashed line is for Hilltop potential with $p = 4$. Red solid line is for $p \to \infty$ in both inverse-Hilltop and Hilltop potentials. For each value of p, the right-side of the red line is excluded because of too much e-foldings. Green line(s) is the bound obtained from the completions of potentials, Eqs. (32), (34) and (33). Blue dot is the prediction of Starobinsky-like potential with $\mu = M_{\rm P}$. *Upper*: $\mu = M_{\rm P}$. *Lower*: $\mu = pM_{\rm P}$ for $p \geq 1$, but $\mu = M_{\rm P}$ for $p < 1$.

THE LOWER BOUND OF r_T: NUMERICAL RESULTS

In order to find a lower bound of r_T, we used the expressions obtained in the previous section and performed a numerical analysis. Also, for comparsion, we used the following completions of potentials:

$$V_{\rm iht} = V_0 \left(1 - 1/2x^p\right)^2 \quad (32)$$

$$V_{\rm st} = V_0 \left(1 - e^{-x}/2\right)^2 \quad (33)$$

$$V_{\rm ht} = V_0 \left(1 - x^p/2\right)^2 \quad (34)$$

The result is shown in Fig. 1. We found that, if $\mu \sim M_{\rm P}$ (upper panels), when $p \gg 4$ in inverse-Hilltop and Hilltop potentials, it is possible to lower-down r_T by many orders of magnitude relative to the current upper bound. However, for $p \leq 4$ which is likely to be the case, in the region of interest the field value is far away from the end point of inflation, and either $r_T \gtrsim \mathcal{O}(10^{-3})$ or n_s is out of the 3-σ bound of observations (in Hilltop). As shown in the lower panels of Fig. 1, if we take μ larger than $M_{\rm P}$, the case of $p \leq 4$ in Hilltop potential can be within the preferred region of observations, but r_T is pushed up to $\mathcal{O}(10^{-3})$. So, we can conlude that, including chaotic monomials and Starobinsky-like potentials, in realistic models (probably) with $p \leq 4$, r_T is lower-bounded at about 10^{-3} or n_s is out-of the 3-σ allowed band. Fig. 1 also shows that, if the decay temperature of the inflaton is low or there is an extra contribution to e-foldings, the lower-bound of r_T is pushed up.

CONCLUSIONS

In this paper, we examined the lower bound of the tensor-to-scalar ratio in large single-field scenarios of inflation. For inflaton field values associated to the relevant cosmological scales, the inflaton potential may be approximated to either chaotic monomial, inverse-Hilltop, Hilltop, or Starobinsky-like potentials in which the leading field-dependent term is $(\phi/\mu)^{\pm p}$ with $0 < p \leq 4$ or $e^{-\phi/\mu}$. We showed that, if the dimensionful scale μ characterizing the end of inflation is Planck scale or larger, which is the case of large single-field scenarios, the tensor-to-scalar ratio is lower-bounded at $\mathcal{O}(10^{-3})$ for the range of the spectral index favored by observations. Therefore, even if it will not be done in the near future, most large single-field inflation models will be probed as experiments reach a r_T at the level of 10^{-3}.

ACKNOWLEDGEMENTS

The authors acknowledge support from the MEC and FEDER (EC) Grants SEV-2014-0398 and FPA2014-54459 and the Generalitat Valenciana under grant

PROME- TEOII/2013/017. G.B. acknowledges partial support from the European Union FP7 ITN INVISIBLES (Marie Curie Actions, PITN-GA-2011-289442).

* Gabriela.Barenboim@uv.es
† Wanil.Park@uv.es

[1] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.01589 [astro-ph.CO].
[2] Q. G. Huang, Phys. Rev. D **91**, no. 12, 123532 (2015) [arXiv:1503.04513 [astro-ph.CO]].
[3] E. W. Kolb and M. S. Turner Addison-Wesley Pub. Co. (1994)
[4] D. H. Lyth and E. D. Stewart, Phys. Rev. Lett. **75**, 201 (1995) [hep-ph/9502417].
[5] D. H. Lyth and E. D. Stewart, Phys. Rev. D **53**, 1784 (1996) [hep-ph/9510204].
[6] S. E. Hong, H. J. Lee, Y. J. Lee, E. D. Stewart and H. Zoe, JCAP **1506**, no. 06, 002 (2015)
[7] D. h. Jeong, K. Kadota, W. I. Park and E. D. Stewart, JHEP **0411**, 046 (2004) [hep-ph/0406136].
[8] S. Kim, W. I. Park and E. D. Stewart, JHEP **0901**, 015 (2009) [arXiv:0807.3607 [hep-ph]].
[9] K. Choi, K. S. Jeong, W. I. Park and C. S. Shin, JCAP **0911**, 018 (2009) [arXiv:0908.2154 [hep-ph]].
[10] W. I. Park, JHEP **1007**, 085 (2010) [arXiv:1004.2326 [hep-ph]].
[11] K. Choi, W. I. Park and C. S. Shin, JCAP **1303**, 011 (2013) [arXiv:1211.3755 [hep-ph]].
[12] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.02114 [astro-ph.CO].